

Trauma in Pregnancy*

Valerie A. Dobiesz | Daniel W. Robinson

PRINCIPLES

Background

Trauma occurs in up to 8% of all pregnancies and is the leading non-obstetric cause of maternal death.¹⁻⁵ The most common causes of injury in pregnancy are motor vehicle collisions (MVCs), interpersonal violence, and falls.^{2,4,6-8} The major determinant of obstetrical outcomes after trauma is the severity of the injury.⁴ Trauma in pregnancy increases the risk of spontaneous abortion, preterm rupture of membranes, preterm birth, uterine rupture, cesarean delivery, placental abruption, and stillbirth.⁴ A significant number of pregnant women admitted to a trauma center do not yet know that they are pregnant.

Commonly used thresholds of fetal viability are an estimated gestational age of 24 to 26 weeks or an estimated fetal weight of 500 g. Only viable fetuses are monitored, because no obstetric intervention will alter the outcome with a previable fetus. Counseling on proper seatbelt and alcohol use and screening for interpersonal violence may help to reduce the morbidity and mortality rates for pregnant patients. Although the essential principles of trauma management remain unchanged in the pregnant patient, a number of special points need to be considered because a gravid uterus does alter the pattern of injury. Pregnancy causes alterations in physiology and anatomy that affect multiple organ systems. Although there are two lives involved, maternal life takes priority and fetal outcomes are directly correlated with early and aggressive maternal resuscitation.⁵

ANATOMY AND PHYSIOLOGY

Anatomic Changes in Pregnancy

The uterus remains an intrapelvic organ until approximately the 12th week of gestation. It reaches the umbilicus by 20 weeks and the costal margins by 34 to 36 weeks. At term, the uterus has often enlarged by 30 cm and has increased fifteen fold in weight, which alters the normal anatomic location and function of multiple structures. The diaphragm progressively rises in pregnancy with compensatory flaring of the ribs, which may predispose to pneumothorax and a faster progression to tension pneumothorax. A thoracostomy done in the third trimester requires that the chest tube be placed one or two interspaces higher than the usual fifth interspace site to allow for diaphragm elevation.

Abdominal viscera are pushed upward by the enlarging uterus and can alter the location of perceived pain. The gravid uterus itself tends to protect abdominal organs from trauma but substantially increases the likelihood of bowel injury from penetrating trauma to the upper abdomen. Conversely, the upward displacement of the bowel makes it less susceptible to blunt trauma.

*The contributors would like to sincerely thank Kriti Bhatia, Hilarie Cranmer, and John D.G. Neufeld for their previous work on this chapter.

The stretching of the abdominal wall as pregnancy nears term modifies the normal response to peritoneal irritation (blunting of muscle guarding and rebound) potentially underestimating the extent and severity of maternal trauma despite significant intra-abdominal bleeding and organ injury.

In the first trimester, the bony pelvis shields the uterus. After the third month, the uterus rises out of the pelvis and becomes vulnerable to direct injury. The bladder is also displaced into the abdominal cavity beyond 12 weeks' gestation, thereby becoming more vulnerable to injury. Like the uterus, the bladder becomes hyperemic, and injury may lead to a marked increase in blood loss compared with similar injury in a nonpregnant patient. Ureteral dilation secondary to smooth muscle relaxation or from compression by the gravid uterus is often found on imaging studies but is not necessarily pathologic. The ligaments of the symphysis pubis and sacroiliac joints are loosened during pregnancy. As a result, a baseline diastasis of the pubic symphysis may exist that can be mistaken for pelvic disruption on a radiograph.

Physiologic Changes

Cardiovascular

The normal cardiovascular changes of pregnancy can alter the clinical presentation and may either mimic or mask the recognition of shock or exacerbate the effects of traumatic hemorrhage (Table 182.1). Blood pressure declines in the first trimester, levels out in the second trimester, and then returns to nonpregnant levels during the third trimester. The decline in systole is small, 2 to 4 mm Hg, whereas diastole falls 5 to 15 mm Hg. Heart rate increases in pregnancy but does not rise by more than 10 to 15 beats per minute above baseline (mean of approximately 90 beats/min).

A major contributor to maternal hypotension is the supine hypotensive syndrome. After 20 weeks' gestation, the enlarging uterus has risen to the level of the inferior vena cava, resulting in compression when the mother is supine. Caval obstruction diminishes cardiac preload, which can decrease cardiac output and systolic blood pressure. In late pregnancy, it is common for the inferior vena cava to become completely occluded when the pregnant patient is supine. Hemodynamic improvement occurs when compression is relieved. In determining whether observed hypotension is related to positioning, the pregnant woman's pelvis can be tilted so that the uterus is displaced from the inferior vena cava (ie, tilt the patient onto her left side unless prevented by other injuries). The uterus can also be manually displaced to the left using two hands and pushing the uterus toward the patient's head. Maintaining a position between 15 and 30 degrees is optimal. Elevating the patient's legs will improve venous return. Inferior vena caval compression can also lower central venous pressure (CVP) in the last two trimesters. Normal CVP during pregnancy is approximately 12 mm Hg. Blood volume gradually increases during pregnancy, starting at 6 to 8 weeks' gestation, to as much

TABLE 182.1

Hemodynamic Changes of Pregnancy (Mean Values)

PARAMETER	NONPREGNANT	TRIMESTER 1	TRIMESTER 2	TRIMESTER 3
Heart rate (beats/min)	70	78	82	85
Systolic blood pressure (mm Hg)	115	112	112	114
Diastolic blood pressure (mm Hg)	70	60	63	70
Cardiac output (L/min)	4.5	4.5	6	6
Central venous pressure (mm Hg)	9.0	7.5	4.0	3.8
Blood volume (mL)	4000	4200	5000	5600
Hematocrit without iron (%)	40	36	33	34
Hematocrit with iron (%)	40	36	34	36
White blood cells (cells/mm ³)	7200	9100	9700	9800

Data from de Swiet M: The cardiovascular system. In Hytten F, Chamberlain G, editors: *Clinical physiology in obstetrics*. Oxford, UK, 1980, Blackwell Scientific Publications, pp 3–42; Colditz RB, Josey WE: Central venous pressure in supine position during normal pregnancy. Comparative determinations during first, second and third trimesters. *Obstet Gynecol* 36:769, 1970; Letsky E: The haematological system. In Hytten RF, Chamberlain G, editors: *Clinical physiology in obstetrics*. Oxford, UK, 1980, Blackwell Scientific Publications, pp 43–78; and Cruikshank DP: Anatomic and physiologic alterations of pregnancy that modify the response to trauma. In Buchsbaum HJ, editor: *Trauma in pregnancy*. Philadelphia, 1979, WB Saunders, pp 21–39.

as 45% above normal, peaking at 32 to 34 weeks' gestation. Blood volumes become increasingly larger for multigravidas and for twin, triplet, and quadruplet gestations. With this increased circulatory reserve, clinical signs of maternal hypotension from acute traumatic bleeding may be delayed. Up to 35% of circulating blood volume may be lost before an injured pregnant patient exhibits signs or symptoms of shock.⁵ By the beginning of the second trimester and throughout the remainder of pregnancy, cardiac output is increased 40% to 6 L/min. Blood flow to the uterus increases from 60 mL/min before pregnancy to 600 mL/min at term. This hyperdynamic state is needed to maintain adequate oxygen delivery to the fetus. Because the mother's total circulating blood volume flows through the uterus every 8 to 11 minutes at term, this can be a major source of blood loss if injured. By the third trimester, there is also marked venous congestion in the pelvis and lower extremities, increasing the potential for hemorrhage from both bony and soft tissue pelvic injuries.

Compression of the lower abdominal venous system by the gravid uterus increases peripheral venous pressure and volume in the legs, creating the potential for brisk blood loss from leg wounds and can exacerbate bleeding from attempts at central venous catheter placement.

Pulmonary

The pregnant woman at term has a significantly reduced oxygen reserve due to a reduction in functional residual capacity caused by diaphragm elevation and an increase in oxygen consumption related to the growing fetus, uterus, and placenta. Mean arterial oxygen tension drops by 29% in pregnant women at term during 60 seconds of apnea compared with 11% in nonpregnant women. Labor further accelerates this decline. In addition, minute ventilation and tidal volume increase, leading to hypocapnia. Therefore a partial pressure of carbon dioxide in the arterial blood (Paco₂) of 35 to 40 mm Hg may indicate inadequate ventilation and impending respiratory decompensation in the pregnant patient. Maternal hypoxia rapidly leads to fetal hypoxia, distress, and possibly demise. There are no contraindications to rapid sequence intubation during pregnancy. Bag-valve-mask ventilation is more difficult in the pregnant patient. The incidence of difficult or failed intubations in obstetric anesthesia is four times higher than in surgical non-obstetric patients.⁸

Gastrointestinal

Gastroesophageal sphincter tone and gastrointestinal motility are decreased in pregnancy, thus increasing the possibility of aspiration in patients with altered level of consciousness, such as during intubation. Early gastric decompression should be performed in these circumstances.

SPECIFIC DISORDERS

Blunt and Penetrating Trauma

Physical examination is unreliable in predicting adverse outcomes in the pregnant woman with blunt trauma. Risk factors predictive of the onset of contractions or preterm labor include gestational age greater than 35 weeks, assaults, and pedestrian collisions. In gravid patients, penetrating trauma of the abdomen increases the likelihood of injury to the bowel, liver, or spleen.

Fetal mortality can be as high as 40% after maternal trauma, with most likely causes of fetal death occurring from placental abruption, maternal shock, and maternal death, in order of decreasing incidence. Risk factors significantly predictive of fetal death include ejection, motorcycle and pedestrian collisions, maternal death, maternal tachycardia, abnormal fetal heart rate, lack of restraints, and an injury severity score greater than 9.

Unbelted or improperly restrained pregnant women are twice as likely to experience excessive maternal bleeding and increased maternal death with fetal death being three times more likely to occur.^{2,4,9,10} For low- to moderate-severity crashes (constituting 95% of all MVCs), proper restraint use, with or without air bag deployment, generally leads to acceptable fetal outcomes. For high-severity crashes, even proper restraint does not improve fetal outcome.

Pregnant crash-test-dummy trials show that improper placement of the lap belt over the pregnant abdomen causes a threefold to fourfold increase in force transmission through the uterus. The lowest force transmission readings through the uterus occur when a three-point seat belt is used properly. For correct position, the lap belt should be placed under the gravid abdomen, snugly over the thighs, with the shoulder harness off to the side of the uterus, between the breasts and over the midline of the clavicle. Women

who receive information on seat belt use during pregnancy from a health care worker are statistically more likely to use seat belts and to use them properly than uninformed controls.

Interpersonal Violence

Physical abuse from husbands or boyfriends compromises a woman's health during pregnancy, as well as her likelihood of carrying a child to term and the health of her newborn. Women experiencing abuse in the year before or during a pregnancy were 40% to 60% more likely than non-abused women to report high blood pressure; vaginal bleeding; severe nausea; kidney or urinary tract infections; and hospitalization during that pregnancy. Abused pregnant women are more likely to deliver preterm, and children of abused pregnant women are more likely to be born underweight. Children born to abused mothers are more likely than other children to require intensive care at birth. Physicians detect only a minority of interpersonal violence cases in pregnant women, which supports the need for routine screening for interpersonal violence in this population.

Falls

Falls become more prevalent after the 20th week of pregnancy and at least one in four pregnant women will fall at least once while pregnant.^{4,6} Protuberance of the abdomen, loosening of pelvic ligaments, strain on the lower back, and fatigability are contributory factors. In a given pregnancy, about 2% of pregnant women sustain repeated direct blows to the abdomen from repetitive falls. Although repeated falls often trigger premature contractions, they seldom result in immediate labor and delivery.

Penetrating Trauma

The gravid uterus affects the injury pattern seen with penetrating trauma to the upper abdomen with the probability of harm to the bowel, liver, or spleen at almost 100%. When the entry site is anterior and below the uterine fundus, visceral injuries are less likely.⁴ Although the enlarging uterus can act as a shield against intra-abdominal injuries in the mother, it makes the fetus more susceptible to injury. A high fetal death rate from penetrating trauma to the uterus has been reported and is lower for maternal injuries above the uterus.

Fetal Injury

There is a high risk of fetal loss in the pregnant trauma patient. Poor fetal outcome is predicted by maternal hypotension and acidosis (hypoxia, lowered pH, lowered bicarbonate) and a fetal heart rate of less than 110 beats/min.⁷ When the mother sustains life-threatening injuries, there is a 40% chance of fetal demise, compared with a less than 2% chance in cases of non-life-threatening maternal injuries. Disseminated intravascular coagulation (DIC), which may be caused by placental products entering the maternal circulation, is a significant predictor of fetal mortality. The American College of Obstetrics and Gynecology recommends a minimum of 2 to 6 hours of fetal monitoring after maternal trauma; cardiotocographic monitoring for a minimum of 4 hours is useful in predicting fetal outcome.⁴

Fatal fetal injuries from blunt trauma are usually the result of intracranial hemorrhage and skull fractures secondary to fractured maternal pelvic bones striking the fetal skull as a result of vertex lie. Pelvic and acetabular fractures during pregnancy are associated with a high maternal (9%) and a higher fetal (38%) mortality rate. Both gunshot wounds and stab wounds to the uterus produce substantial morbidity and mortality to the fetus.

Placental Injury

The leading cause of fetal death after blunt trauma is placental abruption.¹¹ Placental separation results when the inelastic placenta shears away from the elastic uterus during sudden deformation of the uterus. Because deceleration forces can be as damaging to the placenta as direct uterine trauma, abruption can occur with little or no external sign of injury to the abdominal wall. Placental abruption inhibits the flow of oxygen to the fetus and causes in utero carbon dioxide (CO₂) accumulation, resulting in hypoxia and acidosis and leads to fetal distress. Sustained uterine contractions induced by intrauterine hemorrhage also inhibit uterine blood flow, further contributing to fetal hypoxia.

The diagnosis of abruption is made on clinical grounds; ultrasonography and the Kleihauer-Betke test are of limited value. Classic clinical findings of abruption are vaginal bleeding, abdominal cramps, uterine tenderness, maternal hypovolemia (up to 2 L of blood can accumulate in the gravid uterus), or a change in the fetal heart rate; but many cases of placental abruption after trauma present without vaginal bleeding.

The most sensitive indicator of placental abruption is fetal distress, which can be detected with prompt fetal monitoring. Increased frequency of contractions is associated with abruption. Ultrasonography has poor sensitivity for detection of placental abruption (24% sensitivity; 96% specificity).² If the abruption bleeds externally, there may be an insufficient quantity to be detected sonographically. Even with significant intrauterine blood accumulation, accurate ultrasonography diagnosis may be difficult because of placental position (ie, posterior) and confounding uterine or placental structural conditions.

Placental abruption is associated with an increased risk of stillbirth (>20 weeks) and preterm delivery (before 37 weeks) even with minor abruption. The extent of placental separation is correlated with the rate of stillbirth.

A trial of expectant management with ongoing maternal and fetal monitoring is appropriate when mother and fetus are stable and with partial placental abruptions of less than 25%. This usually applies to fetuses of less than 32 weeks' gestation in which the likelihood of morbidity and mortality associated with prematurity makes delivery management risky. Expectant care in stable patients may allow further fetal maturation and improved outcome. An immediate cesarean section should be available in case of fetal distress from further placental separation. After 32 weeks' gestation, the risk of further placental separation outweighs the benefits of further fetal maturation, so intervention is indicated.

Women with placental abruption are more likely to have coagulopathies than those without abruption. The injured placenta can release thromboplastin into the maternal circulation, resulting in DIC, whereas the damaged uterus can disperse plasminogen activator and trigger fibrinolysis. The precipitation of DIC is directly related to the degree of placental separation. Severe clotting disorders rarely occur unless separation of the placenta is significant enough to result in fetal demise.

Uterine Injury

The most common obstetric problem caused by maternal trauma is uterine contractions. Myometrial and decidual cells, irritated by contusion or placental separation, release prostaglandins that stimulate uterine contractions. Progression to labor depends on the extent of uterine damage, the amount of prostaglandins released, and the gestational age of the pregnancy. The routine use of tocolytics for premature labor has come under question because most contractions stop spontaneously. Contractions that are not self-limited are often induced by some pathologic condition, such as underlying placental abruption, which is a contraindication to

tocolytic therapy. Older studies describe this risk as relative and have used tocolysis successfully with careful evaluation and intensive monitoring to continue the pregnancy and enhance fetal maturity. The option to use tocolytics ends when cervical dilation reaches 4 cm.

Uterine rupture is a rare event. It is most often caused by severe vehicular crashes in which pelvic fractures strike directly against the uterus. Uterine rupture may occur from stab wounds and gunshot injuries, but this is rare. Maternal shock, abdominal pain, easily palpable fetal anatomy caused by extrusion into the abdomen, and fetal demise are typical findings on examination. Diagnosing uterine rupture can be difficult. A fractured liver or spleen can produce similar signs and symptoms of peritoneal irritation, hemoperitoneum, and unstable vital signs. Optimal treatment, between suturing the tear or performing a hysterectomy, depends on the extent of uterus and uterine vessel tears and the importance of future childbearing.

DIAGNOSTIC TESTING

All women of childbearing age presenting with trauma should be assessed for possible pregnancy.

CHANGES IN LABORATORY VALUES WITH PREGNANCY

Increases in plasma volume and red blood cells cause physiologic anemia of pregnancy (hematocrit 32% to 34% by the 32nd to 34th week). Despite the lower hematocrit, there is an overall increase in oxygen-carrying capacity because of an increased total red blood cell mass. Placental progesterone directly stimulates the medullary respiratory center, producing a lower PaCO₂ (30 mm Hg) from the second trimester until term. The subsequent compensatory lowering of serum bicarbonate slightly reduces blood-buffering capacity during conditions of physiologic stress. A PaCO₂ of 40 mm Hg in the latter half of pregnancy reflects inadequate ventilation and potential respiratory acidosis that could precipitate fetal distress.

Electrocardiographic changes include a left-axis shift averaging 15 degrees, caused by diaphragm elevation. Consequently, flattened T waves or Q waves in leads III and augmented voltage unipolar left limb lead may be seen.

Laboratory

Laboratory tests for a pregnant patient with trauma should include a blood type with Rh status, arterial blood gas, and coagulation studies. Patients who appear to be stable but have a low serum bicarbonate level may have occult maternal shock. Interpretation of bicarbonate results requires consideration of the physiologic changes that occur in the later stages of pregnancy as a result of respiratory alkalosis. Coagulation studies are important in directing management of patients with multisystem trauma or when the diagnosis of placental abruption is considered.

Kleihauer-Betke Test and Fetomaternal Hemorrhage

Fetomaternal hemorrhage (FMH), the transplacental bleeding of fetal blood into the normally separate maternal circulation, is a unique complication of pregnancy. MVCs, anterior placental location, and uterine tenderness are associated with an increased risk of FMH. Massive fetomaternal transplacental hemorrhage causes alloimmunization in Rh incompatibility but also endangers the fetus by causing severe fetal anemia, fetal distress, and possible exsanguination. ABO incompatibility causes less severe disease.

In theory, it is possible that trauma can result in FMH as early as the fourth week of gestation, when the fetal and placental circulations first form. In practice, FMH is usually of more concern after 12 weeks' gestation, when the uterus rises above the pelvis and becomes susceptible to direct trauma.

The Kleihauer-Betke test quantifies the amount of FMH. Most laboratories screen for FMH of 5 mL or more, even though the amount of FMH sufficient to sensitize most Rh-negative women is well below this 5-mL sensitivity level. Therefore it is advisable that all Rh-negative mothers who have a history of abdominal trauma receive one prophylactic dose of Rhesus immune globulin (RhIG) within 72 hours of injury. Trauma patients at risk for massive FMH will have major injuries or abnormal obstetric findings, such as uterine tenderness, contractions, or vaginal bleeding. Rarely, the amount of FMH will exceed that covered by the maximum RhIG dose (300 µg). Because RhIG can effectively prevent Rh isoimmunization when administered as late as 72 hours after antigenic exposure, the results of the Kleihauer-Betke test are not immediately needed in the emergency department (ED).

Radiography

Plain Radiographs

Adverse effects to the fetus are unlikely if radiation exposure is less than 50 mGy.¹² Less than 1% of trauma patients are exposed to more than 30 mGy. Sensitivity to radiation is greatest during intrauterine development when the embryo undergoes organogenesis in weeks 2 to 15.¹² However, the risk to the fetus of a 10 mGy exposure is thousands of times smaller than the spontaneous risks of malformations, abortions, or genetic disease. Intrauterine exposure to 50 mGy does not appear to cause a significant increase in congenital malformations, intrauterine growth retardation, or miscarriage but is associated with a 2% increase in the risk of cancer.¹² Pathologic conditions more readily appear with intrauterine radiation doses of 150 mGy or greater (at this point, a therapeutic abortion may be considered).¹²

Providing information on radiation exposure from diagnostic radiographs is difficult. Fetal dose from computed tomography (CT) scans depends on many factors, but the type of equipment used, the abdominal girth of the mother, and the fetal distance from the maternal skin are important factors.¹² Diagnostic radiographic studies should be performed with regard for fetal protection, but necessary diagnostic studies of the traumatized pregnant patient should not be withheld out of concern for fetal radiation exposure.¹² When appropriate, fetal irradiation should be minimized by limiting the scope of the examination and using technical means, such as shielding and collimation. [Table 182.2](#) provides estimated radiation doses from various types of examinations.

Ultrasonography

Ultrasonography is the best modality for simultaneous assessment of both the mother and the fetus. In the pregnant patient, it is most useful in detecting major abdominal injury, establishing fetal well-being or demise, gestational age, and placental location.^{4,13} However, ultrasonography has low sensitivity (24%) but high specificity (96%) for placental abruption.^{2,4} Ultrasonography can obviate more hazardous tests, such as CT, cystography, and diagnostic peritoneal lavage (DPL) when a pregnant trauma patient requires an objective evaluation of the abdomen. Limitations in accuracy include operator experience, patient obesity, the presence of subcutaneous air, and a history of multiple abdominal surgeries. If ultrasound findings are equivocal and the patient is hemodynamically unstable, DPL performed in an open

TABLE 182.2

Estimated Fetal Radiation Dose From Conventional Radiographic and Computed Tomography Examination

IMAGING STUDY	ESTIMATED FETAL DOSE (mGy)*
RADIOGRAPHY	
Cervical spine (AP, lateral)	<0.001
Extremities	<0.001
Chest (PA, lateral)	0.002
Thoracic spine	0.003
Abdomen (AP) (21-cm patient thickness)	1
Abdomen (AP) (33-cm patient thickness)	3
Lumbar spine (AP, lateral)	1
COMPUTED TOMOGRAPHY	
Head	0
Chest (routine)	0.2
Chest (pulmonary embolism protocol)	0.2
Abdomen	4
Abdomen and pelvis	25
CT angiography of the aorta	34
CT angiography of the coronary arteries	0.1

*The naturally occurring background radiation dose during pregnancy is 0.5 to 0.1 mGy.

AP, Anteroposterior; CT, computed tomography; PA, posteroanterior.

From Raptis CA, Mellnick VM, Raptis DA, et al: Imaging of trauma in the pregnant patient. *Radiographics* 34(3):748–763, 2014.

and supra-uterine fashion is indicated. Diagnostic peritoneal aspiration (DPA) may be considered in conjunction with a trauma surge; however, this procedure also poses risks.

Computed Tomography and Magnetic Resonance Imaging Scans

CT and, increasingly, magnetic resonance imaging (MRI) studies are used in evaluating abdominal trauma in pregnancy. If ultrasonography is indeterminate and the patient's condition is stable, CT and MRI can identify specific organ damage. They are particularly useful in assessing penetrating wounds of the flank and back. CT can miss diaphragm and bowel injuries. Both of these studies carry the risk of moving the patient from the closely monitored environment of the ED to a distant radiography suite.

Radiation from CT is a concern in the pregnant trauma patient. However, with shielding, fetal exposure from head and chest CT scans can be kept below an acceptable 1-rad limit. CT of the abdomen can be done with 4 mGy of exposure to the fetus.¹⁴ Abdomen and pelvic CT produces about 25 mGy of radiation to the fetus, which is well below the 50 mGy level, where a 2% increase in risk of cancer is seen without evidence of malformation to the fetus.^{12,14} Radiation exposure ultimately depends on the patient, scanner, and technique used in performing the study (see Table 182.2). MRI scanners use no radiation and have not been associated with significant fetal disease or disability.

SPECIAL PROCEDURES

Diagnostic Peritoneal Lavage

In unstable trauma patients with equivocal or negative findings on ultrasonography, DPL can be done in any trimester by an open technique above the uterus after placement of a nasogastric tube and a Foley catheter. The gravid uterus, in the later trimesters, makes the procedure more risky and technically challenging. In blunt trauma, the gravid uterus does not compartmentalize intraperitoneal hemorrhage and does not reduce the accuracy of DPL for identifying patients who need operative intervention for intra-abdominal bleeding. DPL is limited in detecting bowel perforations and does not assess retroperitoneal and intrauterine pathology.

MANAGEMENT

Depending on mechanism, maternal condition, and gestational age, the emergency clinician should consider early notification or consultation with an obstetrician, neonatologist, or pediatrician (or all three). A fetal monitor, portable ultrasound, and neonatal resuscitation equipment should be immediately available. Tetanus toxoid and immune globulin have no detrimental effect on the fetus. The World Health Organization (WHO) specifically recommends vaccination during pregnancy.¹⁵ To prevent alloimmunization of an Rh-negative mother, administer one 50- μ g dose of RhIG in the first trimester. It is sufficient because total fetal blood volume is only 4.2 mL by 12 weeks' gestation, and a 50 μ g-dose covers 5 mL of bleeding. During the second and third trimesters, a 300- μ g dose of RhIG is given, which protects against 30 mL of FMH. Beyond 16 weeks' gestation, the total fetal blood volume reaches 30 mL or more. Massive FMH likely exceeds the efficacy of one 300- μ g dose of RhIG, so the Kleihauer-Betke test can be used to guide effective dosing.

Maternal Resuscitation

Primary Survey

The primary survey focuses on the mother. However, because two patients are present, it is reasonable also to gather preliminary information about the fetus at this time (Fig. 182.1).

Airway and Breathing. Oxygen therapy should be instituted early. Because of reduced oxygen reserve and increased oxygen consumption, the traumatized pregnant woman can quickly become hypoxic. The fetus is very vulnerable to any reduction in oxygen delivery. Supplemental oxygen should be continued throughout maternal resuscitation and evaluation.

A secure airway enables proper oxygenation and negates the higher risk of aspiration in pregnancy. Rapid sequence intubation is recommended. Pre-oxygenation of the pregnant patient is described in Chapter 1. Mechanical ventilation settings need to be adjusted for increased tidal volumes and respiratory alkalosis, which is consistent with the physiologic $Paco_2$ of 30 mm Hg in the last stage of pregnancy.

Circulation. Intravenous access above the diaphragm is preferred.⁸ Maternal blood pressure and heart rate are not consistently reliable predictors of fetal and maternal hemodynamic stability. Due to an expanded circulating volume, the mother can hemorrhage but not show early signs of hypotension. Uterine blood flow is markedly reduced when maternal circulation is compromised. As a result, after an acute blood loss, uterine blood flow can be substantially decreased while maternal blood




Fig. 182.1. Decision-making algorithm in emergency obstetric care. C-section, Cesarean section; FHT, fetal heart tone; US, ultrasonography.

pressure remains normal. Consequently, the pregnant woman with borderline hemodynamic stability probably already has a jeopardized fetus. When traditional signs of shock appear, fetal compromise may be far advanced. Vasopressors should be avoided, because they produce fetal distress by further decreasing uterine blood flow.

Beyond 20 weeks' gestation, tilting the patient to approximately 30 degrees to the left on a backboard or elevating the right hip reduces the compression on the inferior vena cava caused by the gravid uterus. Tilting to the right is less effective in removing the uterus from the inferior vena cava; manually displacing the uterus upward and leftward is more effective.

For severe injuries, a CVP line is helpful in assessing cardiac preload. CVP pressures decline as pregnancy progresses because of inferior vena caval compression by the gravid uterus. Therefore correction to nonpregnant normal pressures may be unnecessary. Instead, it is more valuable to focus on trends of how the CVP responds to fluid resuscitation. A Foley catheter for measuring urine output provides further information on circulatory volume status.

With trauma in pregnancy, the primary survey is modified to assess uterine size and the presence of fetal heart tones if the patient is severely injured. Otherwise, this assessment belongs in the secondary survey. Uterine size, measured from the symphysis pubis to the fundus, is the quickest means of estimating gestational age. This distance in centimeters equals the gestational age in weeks (eg, 24 cm = 24 weeks), which allows some early indication of fetal viability if delivery is necessary (Fig. 182.2). Usually, 24 to 26 weeks is used as the cutoff point for fetal viability (Table 182.3). As a rough guide, the fetus is potentially viable when the dome of the uterus extends beyond the umbilicus. Fetal heart tones can be detected by auscultation at 20 weeks' gestation or by Doppler probe at 10 to 14 weeks. If either the uterus is less than 24 cm in size or fetal heart tones are absent, the pregnancy is probably too early to be viable, and treatment is directed solely at the mother.




Fig. 182.2. Uterine size at different weeks of gestation. (From Kravis TC, Warner CG, editors: Emergency medicine: a comprehensive review. Rockville, MD, 1979, Aspen Publishers.)

TABLE 182.3

Fetal Viability

WEEKS OF GESTATION	6-MONTH SURVIVAL (%)	SURVIVAL WITH NO SEVERE ABNORMALITIES (%)
22	0	0
23	15	2
24	56	21
25	79	69

Data from Morris JA Jr, et al: Infant survival after cesarean section for trauma. Ann Surg 223:481, 1996.

Secondary Survey

The secondary survey involves a detailed examination of the patient but is also modified to gather additional information about the maternal abdomen and the fetus. Physical examination of the abdomen, frequently unreliable in the nonpregnant patient, is even more inaccurate with changing organ position, abdominal wall stretching in advancing pregnancy, and uterine contraction pains. Still, valuable information can be gathered about uterine tenderness, contraction frequency, and vaginal bleeding.

Pelvic examination begins with sterile speculum examination to allow direct visualization to enable detection of possible trauma in the genital tract, the degree of cervical dilation, and the source of any observed vaginal fluid. Vaginal bleeding suggests placental abruption, and a watery discharge suggests rupture of the membranes. If a vaginal fluid sample placed on a slide dries and crystallizes in a ferning pattern, it is amniotic fluid and not urine. Cervical cultures for group B streptococci, *Neisseria gonorrhoeae*, and *Chlamydia* should be performed if there is evidence of amniotic fluid leak. Bimanual examination should be limited to assessing for pelvic bone injury or progression of advanced labor. If the mechanism of injury is significant enough and the fetus is judged

to be viable, early involvement of an obstetrician may enhance the fetal outcome.

Fetal Evaluation. Fetal evaluation in the secondary survey focuses on the fetal heart rate and detection of fetal movement. When the presence of fetal heart tones has been confirmed, intermittent monitoring of fetal heart rate is sufficient for the previsible fetus. If the fetus is viable (ie, 24 weeks or more), continuous external monitoring initiated quickly and maintained throughout all diagnostic and therapeutic procedures may be useful in directing management. Such monitoring can also benefit the mother, because fetal hemodynamics are more sensitive to decreases in maternal blood flow and oxygenation than are most measures of the mother. Fetal distress can be a sign of occult maternal distress. However, fetal distress and even demise can occur with seemingly minor maternal trauma. Signs of fetal distress include an abnormal baseline heart rate, decreased variability of heart rate, and fetal decelerations after contractions.

The normal fetal heart rate ranges from 120 to 160 beats/min; rates outside or trending toward these limits are ominous. Heart rate variability has two components. Beat-to-beat variability measures autonomic nervous function, whereas long-term variability indicates fetal activity. Heart rate variability increases with gestational age. The loss of beat-to-beat and long-term variability warns of fetal central nervous system depression and reduced fetal movement caused by fetal distress (Fig. 182.3).

Late decelerations are an indication of fetal hypoxia. These decelerations are relatively small in amplitude and occur after the peak or conclusion of a uterine contraction. By comparison, early decelerations are larger, occur with the contraction, and recover to baseline immediately after the contraction. Early decelerations may be vagally mediated when uterine contractions squeeze the fetal head, stretch the neck, or compress the umbilical cord. Variable decelerations are large, occur at any time, and are possibly caused by umbilical cord compression (Fig. 182.4).

Mother Stable, Fetus Stable

Minor trauma does not necessarily exempt the fetus from significant injury. It is estimated that up to 3% of all minor trauma

results in fetal loss, typically from placental abruption. Therefore, once the traumatized mother is stabilized, the focus of care is directed toward the fetus. For the viable fetus (greater than 24 weeks' gestation), monitoring is the next step. Continuous monitoring maintained throughout all diagnostic and therapeutic actions is advisable. Because direct impact is not necessary for fetoplacental pathology to occur, the traumatized pregnant woman with no obvious abdominal injury still benefits from monitoring.

The recommended 4 hours of cardiocotographic observation of the viable fetus is extended to 24 hours if at any time during the first 4 hours there are more than three uterine contractions per hour, uterine tenderness persists, results on a fetal monitor strip are worrisome, vaginal bleeding occurs, the membranes rupture, or any serious maternal injury is present. Most cases of placental abruption after maternal trauma are detected within the first 4 hours of monitoring.

On discharge from the hospital, the pregnant woman should be instructed to record fetal movements during the next week. If fewer than four movements per monitored hour are noted, the patient should see her obstetrician immediately and a nonstress test is warranted. The occurrence of preterm labor, membrane rupture, vaginal bleeding, or uterine pain also necessitates prompt reevaluation. Serial ultrasound and fetal heart rate tests on viable fetuses a few days after maternal trauma and periodically throughout the remaining portion of the pregnancy are helpful in monitoring fetal well-being.

Mother Stable, Fetus Unstable

Fetal death rates after maternal trauma are three to nine times higher than maternal death rates.⁵ If a viable fetus remains in distress despite optimization of maternal physiology, cesarean section should be considered.

Although fetal viability is first reached at 24 weeks, the ultimate determinant of the age of fetal viability is the level of neonatal care provided by the intensive care nursery unit in each hospital or accessible regional facility. Determining gestational age for fetuses of less than 29 weeks may be difficult. Emergency decisions on fetal viability are therefore made on the basis of the best ultrasonography and gestational age information available.




Fig. 182.3. Types of fetal heart rate variability. bpm, Beats per minute; FHR, fetal heart rate; UA, uterine activity.




Fig. 182.4. Types of fetal heart rate decelerations. *bpm*, Beats per minute; *FHR*, fetal heart rate; *UA*, uterine activity.

The presence of fetal heart tones is an important survival marker for fetuses about to undergo emergency cesarean section. The fetal survival rate is zero if there are no fetal heart tones present when emergency cesarean section commences. If fetal heart tones are present and the gestational age is 26 weeks or more, then infant survival rate may be as high as 75%.

Besides fetal distress, other reasons for a cesarean section include uterine rupture, placental rupture with significant vaginal bleeding, fetal malpresentation during premature labor, and situations in which the uterus mechanically limits maternal repair. Fetal demise without any of the aforementioned conditions is not an indication for cesarean section, because most will pass spontaneously within 1 week.

Mother Unstable, Fetus Unstable

If the mother's condition is critical, primary repair of her wounds is the best course. This may apply even when the fetus is in distress, because a critically ill mother may not be able to withstand an additional operative procedure such as cesarean section, which prolongs laparotomy time and likely substantially increases blood loss. The best initial action on behalf of the fetus is early and aggressive restoration of normal maternal physiology. If it is felt that the unstable mother can tolerate an emergency cesarean section, it should be considered for the distressed, viable fetus.

As with nonpregnant patients, operative intervention for blunt trauma and above-the-uterus stab wounds is dictated by clinical findings and diagnostic testing results. Above-the-uterus intraperitoneal gunshot wounds require exploration. In situations of severe maternal hemorrhage, massive transfusion protocols should be initiated with fresh frozen plasma, platelets, and red blood cells in a 1:1:1 ratio to lower the rate of coagulopathy and improve survival.⁴ There is little evidence to support a definitive management strategy for penetrating trauma to the gravid uterus. In situations of a hemodynamically stable mother, expectant management has been recommended. However, no prospective study has verified this. Damage to the uterus alone can be quite devastating because of its increased circulation. Without exploration, it is impossible to know the occurrence, size, or depth of

uterine penetration, and there are no guidelines indicating whether a uterine wound can be left unsutured without incurring an increased risk of infection or delayed uterine rupture. Laparotomy or laparoscopy seems to be the safest means of managing penetrating uterine wounds, because missed maternal injuries can quickly compromise the fragile fetus.

Defibrillation

Electrical flow that bypasses the fetus has little effect on the pregnancy. Maternal elective and emergent cardioversion has been performed safely for cardiac dysrhythmias in all three stages of pregnancy. Energies up to 300 Joules have been used without affecting the fetus or inducing premature labor.¹⁶ Although the amount of energy reaching the fetal heart is thought to be small, it is advisable to monitor the fetal heart during maternal cardioversion.

Perimortem Cesarean Section

Restoration of maternal and thus fetal circulation is the optimal goal with maternal hemodynamic instability. However, extended and exclusive attention to the mother in cardiopulmonary arrest may prevent recovery of a potentially viable fetus. During maternal resuscitation, adequate oxygenation, fluid loading, and a 30-degree left tilting position or manual displacement of the gravid uterus, may improve maternal circulation. If there is no response to advanced cardiac life support, a decision for perimortem cesarean section, also referred to as *resuscitative hysterotomy*, should be made within 4 minutes.^{2,17,18} Perimortem cesarean section in the ED should be considered only if uterine size exceeds the umbilicus. Time since maternal circulation ceased is the critical factor in fetal outcome. Delivery increases venous return and cardiac output by 25% to 30% and may lead to survival benefit for mother.^{8,19} Published reports support but fall short of proving that perimortem cesarean delivery should be initiated within 4 minutes of the onset of maternal cardiac arrest and no return of spontaneous circulation.^{8,17,19-22} Beyond 20 minutes, there is virtually never survival or favorable neurologic outcome for either mother or fetus.²

In the event of maternal cardiopulmonary arrest, perimortem cesarean section may be indicated. The most experienced physician available should perform the procedure as cardiopulmonary resuscitation (CPR) is continuing. A midline vertical incision is made from the epigastrium to the symphysis pubis. The uterus is then entered with a midline vertical incision. If necessary, the placenta is incised to reach the fetus; once the fetus has been delivered, the cord is clamped and cut. Maternal revival after delivery of the fetus has been reported in a few perimortem circumstances, presumably because vena caval compression is relieved.

DISPOSITION

Any pregnant woman at 24 or more weeks of gestation who has sustained blunt trauma should undergo at least 4 hours of fetal monitoring even if she looks well. In general, pregnant women who sustain minor trauma have a favorable pregnancy outcome.³ Other admission and operative criteria are similar for pregnant and nonpregnant trauma patients. The emergency clinician should consider the stability of the mother and the viability of the growing fetus when making management and disposition decisions.

KEY CONCEPTS

- Management of life- and limb-threatening injury in the mother comes first. Saving the mother provides the best chance of saving the baby.
- Even in the stable pregnant trauma patient, the fetus is at increased risk of morbidity and mortality.
- The fetus is viable at 24 weeks' gestation. This usually corresponds to when the fundus is at or above the umbilicus.
- Alterations in anatomy and physiology that occur during pregnancy may alter clinical findings related to blood loss and may mask injuries, making methodical assessment essential.
- Stable pregnancies with a viable fetus should be monitored continuously for a minimum of 4 hours after trauma.
- Keeping the mother tilted 30 degrees to the left or manually displacing the uterus may alleviate hypotension and improve perfusion for the mother and fetus.
- Perimortem cesarean section should be initiated for a viable fetus within 4 minutes of no spontaneous return of circulation.
- The use of ionizing radiation to the pregnant patient, including CT and plain radiography, should be minimized, but imaging should not be withheld if it may provide significant diagnostic information. Often, ultrasound, MRI, or a period of observation can preclude the need for ionizing radiation.

The references for this chapter can be found online by accessing the accompanying Expert Consult website.

REFERENCES

- Tinker SC, Reefhuis J, Dellinger AM, et al: Epidemiology of maternal injuries during pregnancy in a population-based study, 1997-2005. *J Womens Health* 19(12):2211-2218, 2010.
- Murphy NJ, Quinlan JD: Trauma in pregnancy: assessment, management, and prevention. *Am Fam Physician* 90(10):717-722, 2014.
- Garmi G, Marjeh M, Salim R: Does minor trauma in pregnancy affect perinatal outcome? *Arch Gynecol Obstet* 290(4):635-641, 2014.
- Mendez-Figuerola H, Dahlke JD, Vrees RA, et al: Trauma in pregnancy: an updated systematic review. *Am J Obstet Gynecol* 209(1):1-10, 2013.
- Battaloglu E, Battaloglu E, Chu J, et al: Obstetrics in trauma. *Trauma* 17:17-23, 2015.
- Petrone P, Talving P, Browder T, et al: Abdominal injuries in pregnancy: a 155-month study at two level 1 trauma centers. *Injury* 42:47-49, 2011.
- John PR, Shiozawa A, Haut ER, et al: An assessment of the impact of pregnancy on trauma mortality. *Surgery* 149:94-98, 2011.
- Einav S, Sela HY, Weiniger CF: Management and outcomes of trauma during pregnancy. *Anesthesiol Clin* 31(1):141-156, 2013.
- Brookfield KF, Gonzalez-Quintero VH, Davis JS, et al: Maternal death in the emergency department from trauma. *Arch Gynecol Obstet* 288(3):507-512, 2013.
- Luley T, Fitzpatrick CB, Grotegut CA, et al: Perinatal implications of motor vehicle accident trauma during pregnancy: identifying populations at risk. *Am J Obstet Gynecol* 208(6):466.e1-466.e5, 2013.
- Barraco RD, Chiu WC, Clancy TV, et al: Practice management guidelines for the diagnosis and management of injury in the pregnant patient: the EAST Practice Management Guidelines Work Group. *J Trauma* 69(1):211-214, 2010.
- Sadro C, Bernstein MP, Kanal KM: Imaging of trauma: part 2, abdominal trauma and pregnancy—a radiologist's guide to doing what is best for the mother and baby. *AJR Am J Roentgenol* 199(6):1207-1219, 2012.
- Puri A, Khadem P, Ahmed S, et al: Imaging of trauma in a pregnant patient. *Semin Ultrasound CT MR* 33(1):37-45, 2012.
- Raptis CA, Mellnick VM, Raptis DA, et al: Imaging of trauma in the pregnant patient. *Radiographics* 34(3):748-763, 2014.
- Keller-Stanislawski B, Englund JA, Kang G, et al: Safety of immunization during pregnancy: a review of the evidence of selected inactivated and live attenuated vaccines. *Vaccine* 32:7057-7064, 2014.
- Trappe HJ: Emergency therapy of maternal and fetal arrhythmias during pregnancy. *J Emerg Trauma Shock* 3(2):153-159, 2010.
- Gatti F, Spagnoli M, Zerbi SM, et al: Out-of-hospital perimortem cesarean section as resuscitative hysterotomy in maternal posttraumatic cardiac arrest. *Case Rep Emerg Med* 2014:121562, 2014.
- Bloomer R, Reid C, Wheatley R: Prehospital resuscitative hysterotomy. *Eur J Emerg Med* 18:241-242, 2011.
- Jeejeebhoy FM, Zelop CM, Windrim R, et al: Management of cardiac arrest in pregnancy: a systematic review. *Resuscitation* 82:801-809, 2011.
- Katz VL: Perimortem cesarean delivery: its role in maternal mortality. *Semin Perinatol* 36(1):68-72, 2012.
- Drukker L, Hants Y, Sharon E, et al: Perimortem cesarean section for maternal and fetal salvage: concise review and protocol. *Acta Obstet Gynecol Scand* 93(10):965-972, 2014.
- Einav S, Kaufman N, Sela HY: Maternal cardiac arrest and perimortem caesarean delivery: evidence or expert-based? *Resuscitation* 83:1191-2000, 2012.

CHAPTER 182: QUESTIONS & ANSWERS

182.1. In cases involving minor trauma to pregnant women, which of the following is a predictor of fetal outcome?

- Abdominal tenderness
- Cardiotocographic monitoring for 4 hours
- Maternal blood count and arterial blood gas results
- Maternal vital signs
- Ultrasonography

Answer: B. For women with mild trauma, fetal outcome is not predicted by maternal vital signs, abdominal tenderness, blood tests, or ultrasound results. Only cardiotocographic monitoring for a minimum of 4 hours is useful to predict fetal outcome.

182.2. Which of the following factors is most concerning in the presentation of a pregnant trauma patient?

- Diastasis of the symphysis pubis
- Electrocardiogram (ECG) findings of Q waves in III and aV_F
- Hematocrit 34% in third trimester
- Hypotension in the third trimester
- Respiratory alkalosis in the third trimester

Answer: D. Blood pressure declines in the first trimester, levels out in the second trimester, and then returns to nonpregnant levels during the third trimester. In pregnancy, minute ventilation increases, leading to hypocapnia. Therefore a partial pressure of arterial carbon dioxide (Paco₂) of 35 to 40 mm Hg may indicate inadequate ventilation and impending respiratory decompensation. The physiologic anemia of pregnancy, resulting from a 48% to 58% increase in plasma volume and only an 18% increase in red blood cells, results in hematocrits of 32% to 34% by gestational age of 32 to 34 weeks. Electrocardiographic changes include a left-axis shift averaging 15 degrees, caused by diaphragm elevation. Consequently, flattened T waves or Q waves in leads III and aV_F may be seen.

182.3. A 26-week gravid woman presents to the emergency department (ED) after a moderate-speed motor vehicle collision (MVC). The patient is without complaints, and her vital signs are as follows: blood pressure,

100/60 mm Hg; heart rate, 100 beats per minute; and respiratory rate, 18 breaths per minute. Ultrasound examination shows good fetal movement, with a fetal heart rate of 150 beats per minute. What is the appropriate disposition for this patient?

- Consult obstetrics for a minimum of 4 hours of cardiotocographic monitoring.
- Consult obstetrics for 1 hour of cardiotocographic monitoring.
- Consult trauma and obstetrics for admission and serial examinations.
- Consult trauma and obstetrics for clearance to discharge patient.
- Discharge the patient with close follow-up with obstetrics.

Answer: A. Placental abruption results when the inelastic placenta shears away from the elastic uterus during sudden deformation of the uterus. Because deceleration forces can be as damaging to the placenta as direct uterine trauma, abruption can occur with little or no external sign of injury to the abdominal wall. For the viable fetus (more than 24 weeks' gestation), monitoring is the next step.

182.4. A 33-year-old pregnant patient complains of abdominal cramping after a trip and fall accident at home. After 4 hours of cardiotocographic monitoring, the patient was admitted to the obstetric service for further evaluation. What was the least likely reason for the admission?

- Late decelerations on fetal monitoring strip
- Leakage of clear fluid from the os
- Maternal pulmonary contusion
- Persistent uterine tenderness
- Two uterine contractions per hour

Answer: E. The recommended 4 hours of cardiotocographic observation of the viable fetus should be extended to 24 hours if at any time during the first 4 hours there are more than three uterine contractions per hour, persistent uterine tenderness, a worrisome fetal monitor strip, vaginal bleeding, rupture of the membranes, or if any serious maternal injury is present.

182.5. A 26-year-old, 30-week gestation woman presents unresponsive with cardiopulmonary resuscitation (CPR) in progress after a high-speed motor vehicle collision (MVC). On ultrasonography, the fetal heart beat is noted at a rate of 130 beats per minute. The patient lost her vital signs 3 minutes before arrival in the emergency department (ED). What is the most appropriate next step in the management of this patient?

- A. Perimortem cesarean section in the operating room by obstetrics
- B. Perimortem cesarean section with horizontal suprapubic incision

- C. Perimortem cesarean section with vertical incision
- D. Thoracotomy with cardiac massage
- E. Thoracotomy with cross-clamping of the aorta

Answer: C. In the event of maternal cardiopulmonary arrest, perimortem cesarean section is indicated. The most experienced physician available should perform the procedure.